



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

Reports of States and yearly and monthly reports of cities of the United States.

GEORGIA—*Columbus*.—Month of November, 1898. Estimated population, 19,303; white, 10,276; colored, 9,027. Total number of deaths, 27 (white, 13, colored, 14), including diphtheria, 1, and 1 from phthisis pulmonalis.

FLORIDA.—Reports to the State board of health for the month of October, 1898, from 45 counties and 5 cities having an aggregate population of 391,422, show a total of 574 deaths, including diphtheria, 2; enteric fever, 116; measles, 3; whooping cough, 1, and 33 from phthisis pulmonalis.

INDIANA.—The reports for the month of November indicate an increase over preceding month, in area of prevalence of tonsilitis, bronchitis, rheumatism, pneumonia, scarlet fever, pleuritis, inflammation of bowels, erysipelas. Decrease indicated in intermittent fever, influenza, diarrhea, and dysentery. Typhoid fever and diphtheria remained unchanged.

Evansville.—Month of November, 1898. Estimated population, 70,000. Total number of deaths, 82, including diphtheria, 1; enteric fever 1, and 4 from phthisis pulmonalis.

MASSACHUSETTS—*Fitchburg*.—Month of November, 1898. Estimated population, 28,392. Total number of deaths, 28, including enteric fever, 1, and 3 from phthisis pulmonalis.

North Attleboro.—Month of November, 1898. Estimated population, 6,725. Total number of deaths, 7, including 1 from phthisis pulmonalis.

Worcester.—Month of October, 1898. Estimated population, 106,202. Total number of deaths, 136, including diphtheria, 5; enteric fever, 2; scarlet fever, 2; whooping cough, 1, and 20 from phthisis pulmonalis.

MICHIGAN.—Reports to the State board of health, Lansing, for the week ended December 3, 1898, from 66 observers, indicate that erysipelas increased and intermittent fever decreased in area of prevalence. Phthisis pulmonalis was reported present at 128, enteric fever at 54, scarlet fever at 35, diphtheria at 30, whooping cough at 14, measles at 10, and smallpox at 2 places—Detroit and Ecorse Township.

MINNESOTA—*Winona*.—Month of November, 1898. Estimated population, 24,000. Total number of deaths, 24, including 2 from phthisis pulmonalis.

NEW YORK—*Yonkers*.—Month of November, 1898. Estimated population, 42,000. Total number of deaths, 50, including 7 from phthisis pulmonalis.

OHIO—*Toledo*.—Month of November, 1898. Estimated population, 151,520. Total number of deaths, 127, including diphtheria, 7; enteric fever, 5, and 10 from phthisis pulmonalis.

PENNSYLVANIA—*Meadville*.—Month of November, 1898. Estimated population, 12,000. Total number of deaths, 4. No deaths from contagious diseases.

SOUTH DAKOTA—*Sioux Falls*.—Month of November, 1898. Estimated population, 11,000. Total number of deaths, 8. No deaths from contagious disease.